

**DEFORMED HEISENBERG ALGEBRA AND FRACTIONAL SPIN
FIELD IN 2+1 DIMENSIONS****Mikhail S. Plyushchay¹**

International Centre for Theoretical Physics, I-34100 Trieste, Italy

ABSTRACT

With the help of the deformed Heisenberg algebra involving Klein operator, we construct the minimal set of linear differential equations for the (2+1)-dimensional relativistic field with arbitrary fractional spin, whose value is defined by the deformation parameter.

MIRAMARE-TRIESTE
September 1993

¹Permanent address: IHEP, Protvino, Moscow Region, 142284 Russia, e-mail: plushchay@mx.ihep.su

The considerable interest to the (2+1)-dimensional particles with fractional spin and statistics (anyons) [1] is conditioned nowadays by their applications to the theory of planar physical phenomena: fractional quantum Hall effect and high- T_c superconductivity [2]. Moreover, anyons attract a great attention due to their relationship with different theoretical fields of research such as conformal field theories, braid groups and deformed theories (see, e.g., refs. [3]–[6]).

From the field-theoretical point of view, such particles can be described in two, possibly related, ways. The first way consists in organizing a statistical interaction of the scalar or fermionic field with the Chern-Simons $U(1)$ gauge field, that changes spin and statistics of the matter field. In this approach, the gauge field can be formally excluded from the theory due to their Lagrange equations (or corresponding Hamiltonian constraints), resulting in the anyonic permutation relations of the redefined matter field [7]. But such an exclusion was intensively criticized, and, till now it is not clear whether the role of the Chern-Simons gauge field is reduced only to the change of the spin and statistics of the matter field, or there is some relic of the statistical gauge field in the theory [8].

Another, less developed way consists in attempting to describe anyons within the group-theoretical approach analogously to the case of integer and half-integer spin fields. The present paper is devoted to further development of this approach, whose program consists in constructing equations for (2+1)-dimensional fractional spin field, subsequent identifying corresponding field action and, finally, in realizing a quantization of the theory to reveal a fractional statistics. Within this approach, there are, in turn, two related possibilities: to use many-valued representations of the $SO(2, 1)$ group, or to work with the infinite-dimensional unitary representations of its universal covering group, $\overline{SO}(2, 1)$ (or $\overline{SL}(2, R)$, isomorphic to it) [9]. Up to now, only the equations were proposed for the fractional spin field within the former possibility [9]–[11], whereas different variants of the equations and field actions were constructed with the use of the unitary infinite-dimensional representations of $\overline{SL}(2, R)$ [9], [12]–[17]. At the same time, the problem of quantizing the theory is still open here. The main difficulty in quantization consists in the infinite component nature of the fractional spin field which is used to describe in a covariant way one-dimensional irreducible representations of the (2+1)-dimensional quantum mechanical Poincaré group $\overline{ISO}(2, 1)$ specified by the values of mass and arbitrary (fixed) spin. Due to this fact, an infinite set of the corresponding Hamiltonian constraints must be present in the theory to exclude an infinite number of the ‘auxiliary’ field degrees of freedom [9]. This infinite set of constraints should appropriately be taken into account. But on the other hand, the infinite component nature of the fractional spin field indicates on the hidden nonlocal nature of the theory, and, therefore, can be considered in favour of the existence of the anyonic spin-statistics relation for the fractional spin fields within the framework of the group-theoretical approach [4, 18].

Relativistic field with arbitrary fractional spin $s = \varepsilon\alpha$, $\alpha > 0$, $\varepsilon \pm 1$, can be described in this approach by the Klein-Gordon equation

$$(P^2 + m^2)\Psi = 0 \tag{1}$$

and (2+1)-dimensional analog of the Majorana equation [19]

$$(PJ - \varepsilon\alpha m)\Psi = 0. \tag{2}$$

Here it is supposed that the field $\Psi = \Psi(x)$ transforms according to the infinite-dimensional unitary irreducible representation (UIR) of the discrete series D_α^+ or D_α^-

of the group $\overline{SL(2, R)}$, and J^μ , $\mu = 0, 1, 2$, are the generators of $\overline{SL(2, R)}$ in the corresponding representation, which satisfy the algebra

$$[J_\mu, J_\nu] = -i\epsilon_{\mu\nu\lambda}J^\lambda \quad (3)$$

and the condition of the irreducibility, $J_\mu J^\mu = -\alpha(\alpha - 1)$ [20]. Eqs. (1) and (2) fix simply the values of the above-mentioned $\overline{ISO(2, 1)}$ Casimir operators, which are the operators of squared mass, $M^2 = -P^2$, and spin, $S = PJ/m$, and moreover, they fix the energy sign: $\text{sign } P^0 = \varepsilon\varepsilon'$, where $\varepsilon' = +1$ and -1 for representations D_α^+ and D_α^- , respectively [9, 12]. These equations are completely independent, unlike, e.g., Dirac and Klein-Gordon equations for the case of spinor field, and, therefore, are not very suitable for constructing the action and quantum theory of the fractional spin field.

In paper [13] Jackiw and Nair constructed linear differential equations for the field with spin $s = \varepsilon(\alpha - 1)$ using a direct product of the vector representation and the UIRs of the discrete series D_α^+ or D_α^- , whereas in ref. [14] more simple set of the Majorana-Dirac equations were proposed for the field with spin $s = \varepsilon(\alpha - 1/2)$ with the use of direct product of the spinor and $D_\alpha^\pm$ representations. In both cases the Klein-Gordon equation was a consequence of the basic linear differential equations, but these equations comprised some subsidiary conditions, which had to be introduced into the corresponding field actions with Lagrange multipliers as some ‘external’ conditions.

In paper [15] Volkov, Sorokin and Tkach proposed the spinor-like set of two linear differential equations for the description of the spin-1/4 particles, whereas in ref. [16] its supersymmetric extension was constructed to describe the supermultiplet with the spin content $s = 1/4$ and $3/4$. It was done with the help of UIRs of the discrete series D_α^+ with $\alpha = 1/4$ or $3/4$. The generators J^μ in these two cases can be represented in the form of operators bilinear in the bosonic oscillator operators $a^\pm$. It is this fact that allows to construct two equations for one spin-1/4 field as a ‘square root’ from the Majorana and Klein-Gordon equations, and to find a spinor representation of this formulation. But, since for the case of arbitrary values α the above-mentioned realization for J^μ in terms of the bosonic oscillator operators $a^\pm$ is impossible, the natural question arises: is it possible to construct in any analogous way the equations for a field with arbitrary fractional spin?

In recent paper [17] the vector set of three linear differential equations for a field with arbitrary fractional spin was constructed. There it was shown that any two of three equations are independent and that the complete vector set of equations is necessary only to have a covariant formulation. Besides, the proposed vector set of dependent equations allowed to demonstrate that only the representations of the discrete series $D_\alpha^\pm$ are appropriate for the description of the fractional spin fields, whereas the possibility to use the infinite dimensional UIRs of the principal and additional continuous series of $\overline{SL(2, R)}$ [20] is excluded.

Therefore, the problem of constructing the minimal covariant set of linear differential equations for the arbitrary fractional spin field is still open, and the present paper is devoted exactly to the solution of it. For the purpose we use here recently proposed realization of the generators J^μ in terms of the operators $a^\pm$ forming the deformed Heisenberg algebra involving the Klein operator [21, 5], and construct the generalization of the spinor-like approach [15] to the case of arbitrary fractional spin field.

So, let us consider the deformed Heisenberg algebra [5, 21, 22]

$$[a^-, a^+] = 1 + \nu Q, \quad Q^2 = 1, \quad Qa^\pm + a^\pm Q = 0, \quad (4)$$

with the self-conjugate Klein operator Q and mutually conjugate operators a^- and a^+ , and introduce the Fock-type vacuum,

$$a^-|0\rangle = 0, \quad Q|0\rangle = \kappa|0\rangle, \quad \langle 0|0\rangle = 1,$$

where $\kappa = +1$ or -1 . Then we get the action of the operator a^+a^- on the states $(a^+)^n|0\rangle$, $n = 0, 1, \dots$,

$$a^+a^-(a^+)^n|0\rangle = \left(n + \frac{1}{2}(1 + (-1)^{n+1})\nu\kappa\right)(a^+)^n|0\rangle.$$

Therefore, the space of unitary representation of algebra (4) is given by the complete set of the normalized vectors

$$|e_n\rangle = \left(\prod_{l=1}^n \left(l + \frac{1}{2}(1 + (-1)^{l+1})\nu\kappa\right)\right)^{-1/2} (a^+)^n|0\rangle, \quad (5)$$

$$\langle e_n|e_k\rangle = \delta_{nk},$$

when $\nu\kappa > -1$. Note that since the number operator is given here by

$$N = a^+a^- - \frac{1}{2}\nu\kappa + \frac{1}{2}\nu Q, \quad N|e_n\rangle = n|e_n\rangle,$$

one can write down the Klein operator in the form

$$Q = \kappa \cos \pi N,$$

and obtain the transcendent equation for defining the operators N , and hence Q , in terms of the deformed creation and annihilation operators:

$$\nu\kappa \cos \pi N = \nu\kappa + 2(N - a^+a^-).$$

With the help of the deformed Heisenberg algebra one can construct the operators J^μ [21],

$$J_0 = \frac{1}{4}(a^+a^- + a^-a^+) = \frac{1}{2}a^+a^- + \frac{1}{4}(1 + \nu Q), \quad J_\pm = J_1 \pm iJ_2 = \frac{1}{2}(a^\pm)^2, \quad (6)$$

which form the algebra (3) of the generators of $\overline{SL(2, R)}$ group. The space of states (5) is divided into two subspaces spanned by the states $|k\rangle_e$ and $|k\rangle_o$:

$$|k\rangle_e \equiv |e_{2k}\rangle, \quad |k\rangle_o \equiv |e_{2k+1}\rangle, \quad Q|k\rangle_{e(o)} = +(-)\kappa|k\rangle_{e(o)}, \quad k = 0, 1, \dots,$$

invariant with respect to the action of operators (6):

$$\begin{aligned} J_0|k\rangle_{e(o)} &= (\alpha_{e(o)} + k)|k\rangle_{e(o)}, \quad k = 0, 1, \dots, \\ J_-|0\rangle_{e(o)} &= 0, \quad J_\pm|k\rangle_{e(o)} \propto |k \pm 1\rangle_{e(o)}, \quad k = 1, \dots \end{aligned} \quad (7)$$

The operator

$$J_\mu J^\mu = -\frac{1}{4}(1 + \nu Q) \left(\frac{1}{4}(1 + \nu Q) - 1 \right)$$

takes the following values on these subspaces:

$$J_\mu J^\mu|k\rangle_{e(o)} = -\alpha_{e(o)}(\alpha_{e(o)} - 1)|k\rangle_{e(o)}, \quad (8)$$

with

$$\alpha_e = \frac{1}{4}(1 + \nu\kappa) > 0, \quad \alpha_o = \alpha_e + \frac{1}{2} > \frac{1}{2}. \quad (9)$$

Relations (7), (8) mean that we have realized the UIRs of the discrete series $D_{\alpha_e}^+$ and $D_{\alpha_o}^+$ of the group $\overline{SL(2, R)}$ on the subspaces spanned by $|k >_e$ and $|k >_o$, respectively. The UIRs of the discrete series $D_{\alpha_{e(o)}}^-$ can be obtained from realization (6) by means of the substitution

$$J_0 \rightarrow -J_0, \quad J_{\pm} \rightarrow -J_{\mp}, \quad (10)$$

and further, for the sake of simplicity, we shall consider only the case of representations D_{α}^+ .

In order to construct the equations for the arbitrary fractional spin field, we introduce the γ -matrices in the Majorana representation,

$$(\gamma^0)_{\alpha}^{\beta} = -(\sigma^2)_{\alpha}^{\beta}, \quad (\gamma^1)_{\alpha}^{\beta} = i(\sigma^1)_{\alpha}^{\beta}, \quad (\gamma^2)_{\alpha}^{\beta} = i(\sigma^3)_{\alpha}^{\beta},$$

which satisfy the relation $\gamma^{\mu}\gamma^{\nu} = -g^{\mu\nu} + i\epsilon^{\mu\nu\lambda}\gamma_{\lambda}$. Here σ^i , $i = 1, 2, 3$, are the Pauli matrices, and raising and lowering the spinor indices is realized by the antisymmetric tensor $\epsilon_{\alpha\beta}$, $\epsilon_{12} = \epsilon^{12} = 1$: $f_{\alpha} = f^{\beta}\epsilon_{\beta\alpha}$, $f^{\alpha} = \epsilon^{\alpha\beta}f_{\beta}$.

Let us consider now the spinor operator

$$L_{\alpha} = \begin{pmatrix} q \\ p \end{pmatrix}, \quad (11)$$

constructed from the operators q and p , which are defined by the operators $a^{\pm}$:

$$a^{\pm} = \frac{q \mp ip}{\sqrt{2}},$$

and in correspondence with eq. (4) satisfy the algebra

$$[q, p] = i(1 + \nu Q), \quad Qq + qQ = Qp + pQ = 0.$$

Taking into account equalities (6), we find that the operators

$$J_{\alpha\beta} \equiv i(J_{\mu}\gamma^{\mu})_{\alpha\beta} = \begin{pmatrix} J_0 + J_1 & -J_2 \\ -J_2 & J_0 - J_1 \end{pmatrix}$$

can be represented in the following form:

$$J_{\alpha\beta} \equiv \frac{1}{4}(L_{\alpha}L_{\beta} + L_{\beta}L_{\alpha}).$$

This equality means that operator (11) is the ‘square root’ operator of the $\overline{SL(2, R)}$ generators (6).

With the help of operator (11), construct the spinor linear differential operator

$$D_{\alpha} = L^{\beta}((P\gamma)_{\beta\alpha} - \varepsilon m\epsilon_{\beta\alpha}), \quad (12)$$

where $P_{\mu} = -i\partial_{\mu}$, $\varepsilon = \pm 1$, and consider the spinor pair of equations

$$D_{\alpha}\Psi = 0. \quad (13)$$

88We suppose here that the field $\Psi = \Psi_e(x)$ or $\Psi = \Psi_o(x)$, is decomposable into the series

$$\Psi_{e(o)}(x) = \sum_{k=0}^{\infty} \psi_{e(o)}^k(x) |k\rangle_{e(o)}. \quad (14)$$

Operator (12) satisfies the relation

$$D^\alpha D_\alpha = L^\alpha L_\alpha (P^2 + m^2),$$

which together with the relation $L^\alpha L_\alpha = -i(1 + \nu Q) \neq 0$ means that the Klein-Gordon equation (1) is the consequence of eqs. (13). Moreover, due to the relation

$$L^\alpha D_\alpha = -4i \left(PJ - \varepsilon m \frac{1}{4} (1 + \nu Q) \right),$$

we conclude that the fields Ψ_e and Ψ_o satisfying eqs. (13), obey also the corresponding equations

$$(PJ - \varepsilon \alpha_e m) \Psi_e = 0, \quad (15)$$

$$\left(PJ - \varepsilon \left(\alpha_o - \frac{1}{2} (1 + \nu \kappa) \right) m \right) \Psi_o = 0. \quad (16)$$

Eq. (15) is nothing else as the Majorana equation (2). Taking into account eq. (1), and passing over to the rest frame $\mathbf{P} = \mathbf{0}$ in the momentum representation, we find with the help of eq. (6) that eq. (15) has the solution of the form $\Psi_e \propto \delta(P^0 - \varepsilon m) \delta(\mathbf{P}) \psi_e^0 |0\rangle_e$, whereas eq. (16) has no nontrivial solution. Therefore, the pair of equations (13) has nontrivial solutions only in the case $\Psi = \Psi_e$, i.e. when $Q\Psi = \kappa\Psi$, describing the field with spin $s = \varepsilon \alpha_e$ and mass m . At $\nu = 0$, i.e. when $[a^-, a^+] = 1$, eqs. (13) in correspondence with eqs. (9), (15), turn into Volkov-Sorokin-Tkach equations for a field Ψ_e with spin $s = 1/4 \cdot \varepsilon$ [15] being analogous to (3+1)-dimensional Dirac positive-energy relativistic wave equations [23].

Thus, we have constructed the spinor-like system of linear differential equations (13) being the independent equations for the field $\Psi = \Psi_e$ with arbitrary fractional spin defined by the deformation parameter: $s = \varepsilon \cdot \frac{1}{4} (1 + \nu \kappa) \neq 0$. Eqs. (13) are in fact the ‘square root’ of the Klein-Gordon and Majorana equations and they can be used as a starting point to construct the quantum theory of the anyons within the group-theoretical approach. In future publications we hope to investigate the problems of constructing the field action leading to equations (13) and quantization of the theory, and, besides, we will consider the supersymmetric extension of the proposed formulation for the arbitrary fractional spin field.

Acknowledgements.

The author thanks Prof. Abdus Salam, the International Atomic Agency and UNESCO for hospitality at the International Centre for Theoretical Physics, Trieste, where the present work was completed.

He is also grateful to A.T. Filippov, P.A. Marchetti, J. Myrheim and M.V. Vasiliev for useful discussions, and to M. Tonin for discussions and hospitality at Dipartimento di Fisica, Università di Padova.

References

- [1] J.M. Leinaas and J. Myrheim, *Nuovo Cim.* **B37** (1977) 1;
F. Wilczek, *Phys. Rev. Lett.* **48** (1982) 1144; **49** (1982) 957
- [2] R. Mackenzie and F. Wilczek, *Intern. J. Mod. Phys.* **A3** (1988) 2827;
F. Wilczek, *Fractional Statistics and Anyon Superconductivity*, (World Scientific, Singapore, 1990);
S. Forte, *Rev. Mod. Phys.* **64** (1992) 193;
K. Lechner and R. Iengo, *Phys. Rep.* **213** (1992) 179
- [3] S. Fubini, *Condensed Matter and High Energy Physics*, preprint CERN-TH-5763/90, 1990; *Mod. Phys. Lett.* **A6** (1991) 347
- [4] J. Fröhlich and P. Marchetti, *Nucl. Phys.* **B356** (1991) 533;
P.A. Marchetti, *Intern. J. Mod. Phys.* **B6** (1992) 2275
- [5] L. Brink, T.H. Hansson and M.A. Vasiliev, *Phys. Lett.* **B311** (1992) 109;
L. Brink, T.H. Hansson, S. Konstein and M.V. Vasiliev, *Nucl. Phys.* **B401** (1993) 591
- [6] A. Lerda and S. Sciuto, *Nucl. Phys.* **401** (1993) 613
- [7] G.W. Semenoff, *Phys. Rev. Lett.* **61** (1988) 517
- [8] P. Gerbert, *Intern. J. Mod. Phys.* **A6** (1991) 173;
A. Lerda, *Anyons*, Lecture Notes in Physics, **m14** (Springer, Berlin, 1992), and references therein
- [9] M.S. Plyushchay, *Intern. J. Mod. Phys.* **A7** (1992) 7045
- [10] M.S. Plyushchay, *Phys. Lett.* **B248** (1990) 107
- [11] S. Forte and T. Jolicoeur, *Nucl. Phys.* **B350** (1991) 589
- [12] M.S. Plyushchay, *Phys. Lett.* **B262** (1991) 71; *Nucl. Phys.* **B362** (1991) 54
- [13] R. Jackiw and V.P. Nair, *Phys. Rev.* **D43** (1991) 1933
- [14] M.S. Plyushchay, *Phys. Lett.* **B273** (1991) 250
- [15] D.V. Volkov, D.P. Sorokin and V.I. Tkach, in *Problems in Modern Quantum Field Theory*, A. Belavin, A. Klimyk and A. Zamolodchikov, eds. (Springer, Berlin, 1989)
- [16] D.P. Sorokin and D.V. Volkov, *(Anti)commuting spinors and supersymmetric dynamics of semions*, preprint IC-92-121, Trieste, 1992; *JETP Letters* **57** (1993) 330
- [17] J.L. Cortés and M.S. Plyushchay, *Linear Differential Equations for a Fractional Spin Field*, DFTUZ preprint 92/24, Zaragoza, 1992
- [18] D. Buchholz and K. Fredenhagen, *Comm. Math. Phys.* **84** (1982) 1;
J. Fröhlich and P. Marchetti, *Lett. Math. Phys.* **16** (1988) 347; *Comm. Math. Phys.* **121** (1989) 177

- [19] E. Majorana, *Nuovo Cim.* **9** (1932) 335;
D.Tz. Stoyanov and I.T. Todorov, *J. Math. Phys.* **9** (1968) 2146
- [20] V. Bargmann, *Ann. Math.* **48** (1947) 568;
D.P. Zhelobenko and A.I. Stern, *Representations of Lie groups* (Nauka, Moscow, 1983) [in Russian];
M.S. Plyushchay, *J. Math. Phys.* **34** (1993) 3954
- [21] M.V. Vasiliev, *Pis'ma JETP* **50** (1989) 344; *Inter. J. Mod. Phys.* **A6** (1991) 1115
- [22] T. Brzezinski, I.L. Egusquiza and A.J. Macfarlane, *Phys. Lett.* **B311** (1993) 202
- [23] P.A.M. Dirac, *Proc. Roy. Soc.* **A322** (1971) 435, **328** (1972) 1572